

Geomagnetic Survey to Explore High-Temperature Geothermal System in Blawan-Ijen, East Java, Indonesia

Yunus Daud^{1,2*}, Syamsu Rosid^{1,2}, Fikri Fahmi¹, Faris Maulana Yunus² and Reza Muflihendri²

¹Master Program in Geothermal Exploration, Graduate Program of Physical Science, Universitas Indonesia, Jakarta, Indonesia

²Study Program of Geophysics Faculty of Mathematics and Natural Science, Universitas Indonesia, Depok, Indonesia

Abstract. Ijen geothermal area is high-temperature geothermal system located in Bondowoso regency, East Java. It is categorized as caldera-hosted geothermal system which is covered by quaternary andesitic volcanic rocks with steep topography at the surrounding. Several surface thermal manifestations are found, such as altered rocks near Mt. Kukusan and a group of Blawan hot springs in the northern part of the caldera. Geomagnetic survey was conducted at 72 stations which is distributed inside the caldera to delineate the existence of hydrothermal activity. Magnetic anomaly was obtained by reducing total magnetic measured on the field by IGRF and diurnal variation. Reduction to pole (RTP) method was applied with geomagnetic inclination of about -32° . In general, the result shows that high magnetic anomaly is distributed at the boundary of study area, while low magnetic anomaly is observed in the centre. The low anomaly indicates demagnetized rock that probably caused by hydrothermal activity. It has a good correlation with surface alteration observed close to Mt. Kukusan as well as high temperature reservoir drilled in the centre of caldera. Accordingly, the low magnetic anomaly also presents the possibility of geothermal reservoir in Ijen geothermal area.

1 Introduction

Ijen Volcanic Complex is located in the Bondowoso, West Java, Indonesia. Expectation of geothermal system appearance in the volcanic complex is indicated by the occurrence of Blawan hot spring in the northern part of caldera rim and steam heated extremely acid sulphate water on Ijen crater as the only surface manifestation in this area. The hot spring express an outflow of the system, while the Ijen crater indicates a young volcano which is uneconomical to be developed. Therefore, it is necessary to do further research to find an appropriate location to be targeted in such challenging hidden geothermal system.

In 2017, magnetic survey had been conducted over Ijen geothermal prospect area at 72 stations. Magnetic method is carried out to measure magnetic field variations in the earth's surface. The studies of magnetic anomalies are often useful for investigating high temperature geothermal systems hosted by Quaternary volcanic rocks [1]. These anomalies represent the demagnetization of reservoir rocks caused by hydrothermal processes due to thermal fluid and rock interactions. In this study, we used a reduce to pole (RTP) technique for imaging total force of magnetic anomalies. RTP transformation delineates the hydrothermally demagnetised rock which is shown as negative anomalies [2]. This paper presents the results of magnetic anomaly, RTP anomaly and the upward-continuation anomaly which were then interpreted to get

a conclusion of the geothermal prospect zone in the Ijen Volcanic Complex.

2 Regional Geology

The Ijen Volcanic Complex (IVC) is located in East Java. The Old Ijen volcano is thought to form around the Pleistocene [3]. The caldera itself is formed by Plinian eruption, ejecting 80 km³ of volcanic material, back to the 0,2-0,05 Ma [4]. Currently, the product is revealed partly at the northern part. Meanwhile, in the southern part, it has been covered by younger volcanoes.

The post-caldera volcanoes are classified into two groups: Caldera Rim Volcanoes and Intra Caldera Volcanoes [5]. Some fault are found in the caldera and extends right in the centre of the caldera, strikes the remains caldera rim in the north and tear down the wall into a steep creek. Here in IVC, several structures that occurred are Blawan fault, Kawahwung fault, Krepekan fault, Cemara-Kukusan fault, Kalipahit-Banyulinu fault, Djampit fault, Rante fault, Pawenan-Blau fault and Kendeng-Merapi fault (Fig.1).

The deformation may caused by tectonic activity, magma refilling in the chamber, or collaboration of those two that reactivate the caldera floor structures. Hot springs in Blawan area indicating that secondary permeability will affect the geothermal manifestation occurrences.

* Corresponding author: ydaud@sci.ui.ac.id




Fig. 2. Magnetic Anomalies over Ijen Geothermal Area

4.3 Upward Continuation

The pattern of negative anomaly found in the centre of the study area is probably associated with the presence of hydrothermal demagnetised rocks of Mt Kukusan. It is supported by the presence of surface alteration observed close to Mt. Kukusan.

The location of the prospect zone is also strengthened by the intersection of several faults; Blawan fault, Cemara-Kukusan fault, and Kawahwuring Fault which could form a high permeable zone.

Fig. 4 shows the upward continuation anomaly map. Generally, the pattern of the high and low magnetic anomaly in the upward continuation map yields the same pattern with the RTP anomaly.

The upward continuation anomaly represents a regional anomaly. As shown in the result, it further strengthens that the negative magnetic anomalies area occurred in the northern part of the study area and extended to the southeast which could become the prospect zone in Blawan Ijen geothermal area.




Fig. 3. The result of Reduce to Pole (RTP) of Magnetic Anomalies of Ijen Geothermal Area

