

Biodecolorization of anthraquinone dyes using immobilised mycelium of *Bjerkandera adusta* CCBAS930

Kamila Rybczyńska-Tkaczyk¹, Teresa Kornilłowicz-Kowalska¹

¹University of Life Sciences in Lublin, Department of Environmental Microbiology, Laboratory of Mycology, Leszczyńskiego 7 Street, 20-069 Lublin, Poland, e-mail: kamila.rybczynska-tkaczyk@up.lublin.pl

Abstract The aim of this study was to characterize the activity of oxidoreductases during biotransformation of 0.01% anthraquinone dyes: Alizarin Blue Black B (ABBB) and Acid Blue 129 (AB129), Carminic Acid (CA), Remazol Brilliant Blue R (RBBR), Acid Green 25 (AG25) and Poly R-478 by immobilized strain of *Bjerkandera adusta* CCBAS 930. Phenolic compounds, phytotoxicity (*Lepidium sativum* L.), biotoxicity were evaluated to determine the toxicity of anthraquinone dyes before and after the treatment with immobilized *B. adusta* CCBAS 930. More than 60% of CA and AB129 were removed by immobilized *B. adusta* CCBAS after 7 days. No secondary products toxic to plants and bacteria were formed during immobilized cultures of *B. adusta* CCBAS 930.

Keywords: white-rot fungi, versatile peroxidase, anthraquinone dyes, immobilization

1. Introduction

Synthetic dyes are widely used in many industries, such as textile, cosmetic, pharmaceutical and food industries. Anthraquinone dyes are most widely used for the dyeing of cotton, wool and silk. Textile industry produces the largest amount of dye effluents. In recent years, with the increase in demand for textile products, the global textile production has increased rapidly and reached about 140.84 million tons in 2012 [1]. Currently, wastewater containing textile effluents are treated using physicochemical methods: ozonation, adsorption and membrane filtration [2]. In recent years biological methods using microorganisms to remove contaminants are gaining in importance, supplementing or presenting an alternative to conventional methods. The biological methods are attractive because of their low costs, environmental safety and common societal acceptance. Particularly promising in this regard are white-rot fungi (*Basidiomycetes*). The most frequently mentioned biological mechanism of contaminant removal by these organisms is biodegradation, involving exploitation of the natural ability of the fungi to synthesize extracellular ligninolytic enzymes. Owing to the low substrate specificity of these enzymes, they can be used for biodegradation of natural and synthetic aromatic compounds that are structurally related to lignin [3]. Decolorization and biodegradation abilities of different *Bjerkandera* spp., including *B. adusta* strains, were widely studied, and basically include azo, anthraquinonic, triphenylmethane and heterocyclic dyes. Studies using the anamorphic stage of *Bjerkandera*

adusta CCBAS 930 have shown that this strain can remove a wide spectrum of aromatic compounds from an aquatic environment. In previous studies the *B. adusta* CCBAS 930 strain has been shown to exhibit decolorizing activity on synthetic dyes with an anthraquinone structure (monoanthraquinones: Carminic Acid and Remazol Brilliant Blue R; polyanthraquinones: Poly-R) [4,5], Alizarin Blue Black B, Acid Blue 129 (data not show), triphenylmethane dyes (Brilliant Green) and heterocyclic dyes (erythrosine) [6]. The use of such immobilized cultures for biodegradation of aromatic compounds has many advantages. Immobilized cells are more stable, durable, and resistant to environmental conditions than conventional cultures [7,8].

The aim of this study was to determine the effectivenesses of anthraquinone dyes: Alizarin Blue Black B (ABBB), Acid Blue 129 (AB129), Carminic Acid (CA), Remazol Brilliant Blue R (RBBR), Acid Green 25 (AG25) and Poly R-478 decolorization and peroxidases production by immobilised mycelium of *B. adusta* CCBAS 930.

2. Materials and Methods

2.1. Strain of *Bjerkandera adusta* CCBAS 930

Anamorphic stage of *B. adusta* CCBAS 930 was isolated from the black earth soil (Pheozems, FAO classification) from a field near Lublin, South-Eastern Poland [9].

1.2. Immobilisation of mycelium of *B. adusta* CCBAS 930

*Corresponding author: kamila.rybczynska-tkaczyk@up.lublin.pl

One hundred ml of a sterile Na-alginate solution (4%) were mixed with 20 g of homogenized mycelia of *B. adusta* CCBAS 930 and its mutants 930-5 and agitated (150 rpm, 20 min). The mixture (10 ml) was extruded using a sterile syringe into 0.2M CaCl₂, thus forming beads with a 3.0-4.0 mm diameter. The beads were allowed to harden in 0.2M CaCl₂ for about 24 h at 4°C. Next, the 0.2M CaCl₂ solution was removed and the beads were washed twice with the addition of distilled water. Immobilised mycelium (Na-alginate beads) of *B. adusta* CCBAS 930 was incubated with 0.01% solutions of anthraquinone dyes (150 rpm, 28°C, 7 days).

2.3. Biochemical analysis of supernatants

Samples were collected every day and following parameters were estimated: the degree of decolorization of anthraquinone dyes using an UV/VIS spectrophotometer [10] content of phenolic compounds [11] and activity of horseradish-type (HRP-like) and versatile (VP). The activity of HRP-like was estimated using 255 µl of 0.01% o-dianisidine ($\epsilon_{460nm}=11,3 \text{ M}^{-1} \text{ cm}^{-1}$) in 0.1 M acetate buffer (pH 5.5) with 0.1 mM H₂O₂. Versatile peroxidase (VP) activity was determined by the oxidation of 20 mM 2,6-dimethoxyphenols (2,6-DMP). Mn-independent activity of VP was estimated by the oxidation of 15 µl of 20 mM 2,6-DMP ($\epsilon_{468} = 49.6 \text{ M}^{-1} \text{ cm}^{-1}$) in 265 µl 50 mM sodium malonate buffer (pH 3.0 and 4.5) with 10 µl of 6 mM H₂O₂. Mn-dependent activity of VP was assayed by oxidation of 15 µl 20 mM 2,6-DMP in 250 µl 50 mM sodium malonate buffer (pH 4.5) in the presence of 10 µl of 6 mM H₂O₂ and 15 µl 0.1 mM MnSO₄. One unit of specific enzyme activity (U mg⁻¹) was defined as the amount of enzyme that oxidized 1 µM of the substrate per minute under defined conditions. The protein concentration was determined according to the Bradford method (1976) [12] using the Protein Assay Kit (BioRad).

2.4. Phytotoxicity assay

The phytotoxicity assay was performed for the untreated medium with 1% molasses and supernatants obtained after 21-day stationary cultures and 7-day immobilized cultures of *B. adusta* CCBAS 930 and its 930-5 mutant. The phytotoxicity assay was carried out to determine root growth inhibition (I) and germination capacity (GI) in *Lepidium sativum*. L. seeds before and after decolorization [13]. Petri dishes were lined with filter paper and, subsequently, 100 seeds and 5 ml of the filtered ($\phi=0.22 \mu\text{m}$) sample were added. The dishes were incubated at room temperature for 72 hours. Distilled water was used as a control.

2.5. Biototoxicity assay using resazurin reduction method

The initial 0.01% dye solution and post-culture fluids from immobilized cultures of *B. adusta* CCBAS 930 strain were tested against Gram-positive *Staphylococcus aureus* ATCC 29737 and Gram-negative *Escherichia coli* ATCC 25922. The strains were provided by ATCC and stored at 4°C. All strains were cultured on nutrient broth (NB) medium at 37°C. Resazurin is a non-toxic water-soluble dye previously applied in bacterial viability studies [14]. This assay is based on detection of the metabolic activity of the cells. The redox dye resazurin (7-hydroxy-3H-phenoxazin-3-one 10-oxide) enters the cell in the oxidized form (blue), where it is converted to a reduced form, resorufin (pink). The reduced and oxidized forms of resazurin can be measured separately with a spectrophotometer and used to determine the reduction capability of cells, which reflects the mitochondrial function and cell viability and shows time- and concentration-dependent cell growth inhibition. Serial twofold dilutions of initial 0.01% dyes solution and post-culture fluids from immobilized cultures of *B. adusta* CCBAS 930 were made with Mueller Hinton Broth (MHB) to yield final concentrations ranging from 10 to 0.62 mg ml⁻¹ (1 - 0.062%) and placed into a 96-well plate. The bacterial suspension (100 µl) prepared from an overnight culture was adjusted to inoculation of 10⁸ CFU mL⁻¹. Then, 100 µL of bacterial culture were added. The wells with MHB or yeast culture were the negative and positive control, respectively. The plates were incubated at 37°C for 48h. Then, 20 µl of a 60-µM resazurin solution in PBS buffer were added to each well. After incubation (2h, 37°C), the viability of cells was monitored by measuring absorbance at 570 nm (reduced) and 600 nm (oxidized) [14] and calculating the resazurin reduction factor (RRF).

2.6. Data analysis

The results were presented as a mean of three repetitions for which standard deviation was calculated.

3. Results

The results showed that the use of immobilised mycelium of *B. adusta* CCBAS 930 significantly reduces the time of decolorization of AB33, AB129, CA and RBBR. After 7 days decolorization degree of CA, AB129, AB33 and RBBR was 75.50%, 65.08%, 56.57% and 25,64%, respectively (Fig. 1).




Fig. 1. Decolorization degree of anthraquinone dyes in immobilized cultures of *B. adusta* CCBAS 930

In immobilised cultures of *B. adusta* CCBAS 930 with 0.01% of anthraquinone dyes activity of all tested peroxidases was detected, but HRP-like and VP was characterized by the highest activity. Activity of HRP-like and VP systematically increased reaching the maximum after 7 days, 35.64 - 80.32 and 50.20 - 106.43 U mg⁻¹ protein, respectively (Fig.2).




Fig. 2. Activities of peroxidases during immobilized cultures of *B. adusta* CCBAS 930 in the presence of anthraquinone dyes

During the immobilised culture of *B. adusta* CCBAS 930 with 0.01% anthraquinone dyes, a systematic decrease in the content of phenolic compounds was observed (50-60%) (Fig. 3).




Fig. 3. Content of phenolic compounds during biotransformation of anthraquinone dyes by immobilized *B. adusta* CCBAS 930

Decolorized post-culture fluids were characterized by lower phyt- and biotoxicity. Decolorized post-culture liquids were characterized by a similar level of inhibition of root growth as control medium initial solution of tested dyes. After biological treatment we observed higher germination index compared to the control solution (Fig.4).




Fig. 4. Root growth inhibition (RGI) and germination index (GI) before (A1, B1) and after decolorization (A2, B2) in cultures of *B. adusta* CCBAS 930

The biotoxicity assay with Gram-positive and Gram-negative *bacteria* demonstrated a similar RRF value for the supernatants after immobilized cultures of *B. adusta* CCBAS 930 and control bacteria cultures. Based on these results, it can be concluded that no toxic products are formed during removal of anthraquinone dyes by immobilized *B. adusta* CCBAS (Fig. 5).




Fig.5. Resazurine reduction factors (RRF) (C,D) of control solutions of 0.01% anthraquinone dyes and decolorized post-culture liquids from immobilized cultures of *B. adusta* CCBAS 930

4. Discussion

White rot fungi have the ability to remove dyes of a complex structure, and immobilization of mycelium increase the efficiency of this process, including reduced time of dyes removal [15]. Compared to conventional cultures, the use of immobilized mycelium allows for more effective removal of synthetic anthraquinone dyes and gives the possibility of using a biological method in various industries [8,16]. In addition, liquids after immobilized cultures do not show biotoxicity or phytotoxicity as opposed to traditional cultures. In conventional cultures, various metabolites are formed during the growth of the fungus in the presence of synthetic dyes. Secondary products may also be formed which may be toxic [17].

Based on the results obtained in this study, it can be concluded that the removal of anthraquinone dyes using immobilized mycelium of *B. adusta* CCBAS 930 is more effective and efficient compared to cultivating this

strain in stationary conditions. After 7 days, removal of RBBR and Poly R-478 dye was observed at 25% and 10%, respectively. Studies of Kornilowicz-Kowalska and Rybczyńska (2012) [4] on removal of 0.01% RBBR and 0.03% Poly-R-478 in liquid cultures of *B. adusta* CCBAS 930 showed only a 5% and 12% decrease in color medium, respectively. After 18 days of culture, total decolorization of 0.01% RBBR was noted. In contrast, complete decolorization of polyanthraquinone dye Poly-R 478 was observed on 30 day of culture. Research of Kornilowicz-Kowalska and Rybczyńska (2014) [5] on the cultivation of strain *B. adusta* CCBAS 930 with 0.01% AC under static conditions showed that after 4 days, the basidiomycete removed more than 72%. This corresponded to a 76.82% reduction in this dye concentration in the medium. The results obtained in this work show that the removal of 0.01% AC in the *B. adusta* CCBAS 930 immobilized culture was not accelerated. In the case of 0.01% AC, the final effect is similar to the results obtained in *B. adusta* CCBAS 930 stationary culture. In the research of L'opez et al. (2006) [18] Poly R-478 and RBBR in a concentration of 200mg L⁻¹, were removed by 31 fungi. In the case of Poly R-478, only 6 strains showed effective, over 90% decolorization after 7 days. The authors also found that RBBR was decolorized only by three strains LFM-4, LFM-19, LFM-22 at 50% after 7 days [18]. The results of this study can confirm the above conclusion because RBBR and Poly R-478 dyes from the five analyzed were the most difficult to biodegrade by the immobilized mycelium *B. adusta* CCBAS 930

Our previous study indicated that in stationary culture of *B. adusta* with the addition of Poly R-478 and RBBR, a decrease in phenolic compounds noted. A decrease in phenolics was observed after 7 and 10 days of fungal growth [4]. In addition, in the optimized cultures of microscopic fungi *H. hematococca* BwIII43, K37 and *T. harzianum* BsIII33 with the addition of 0.03% ABBB dye found that the content of phenolic compounds decreased by 70-76% and 30% respectively after 2 weeks [19]. A 43% drop in phenolic compounds after 7 days of immobilized cultures of *B. adusta* CCBAS 930 with 0.01% ABBB allows to conclude that the immobilization affects the acceleration of the decrease in phenolic compounds content during ABBB biodegradation. On this basis, the effectiveness of immobilization can be determined compared to stationary culture.

The results of this study show that anthraquinone dyes are removed by the immobilized mycelium *B. adusta* CCBAS 930 using extracellular peroxidases: HRP-like and (VP). The results of the obtained research are consistent with the previous ones, which describe the decolorization of anthraquinone dyes by fungal peroxidases [4,5,10]. The results of the research by Rybczyńska-Tkaczyk et al. (2018) [10] showed that other fungi like *H. haematococca* BwIII43, K37 and *T. harzianum* BsIII33 due to the possibility of synthesizing extracellular peroxidases are able to remove 0.03% ABBB monoanthraquinone dye.

This study on the immobilized mycelium *B. adusta* CCBAS 930 also indicates that the test strain is able to decolorize 0.01% ABBB at 56.5% in 7 days using extracellular peroxidases VP and HRP-like. Our earlier studies show that the stationary cultures of *H. hematococca* BwIII43, K37 and *T. harzianum* BsIII33 with the addition of 0.03% ABBB showed the activity of enzymes such as laccase, LiP, MnP and HRP-like. In the cultures of these mushrooms, the highest HRP-like peroxidase activity was recorded in the first week and it was 106,63–109,47 U*mg⁻¹ [19]. In cultures of *B. adusta* CCBAS 930 with 0.01% Poly R-478 and AC, maximum HRP-like peroxidase activity was observed on 18 and 21 days - 33,81 mU*ml⁻¹ or 13,27 mU*ml⁻¹ respectively [5]. Tinoco et al. (2007) [20] studied the removal of industrial dyes by VP peroxidase from *B. adusta* UAMH 8258 and found significant differences in the dyeing of these dyes. Moreira et al. (2006) [21] isolated VP from *Bjerkandera* sp B33/3, which effectively removed RBBR dye. According to the authors, VP peroxidase from B33/3 strain has unique properties. However, they show differences between VP peroxidases synthesized by different *Bjerkander* strains.

An important aspect of biodegradation is also the toxicity of anthraquinone dyes. Most often, the decrease in the toxicity of synthetic dyes is measured using bacteria and *L. sativum* L. seeds [10,13]. The bio- and phytotoxicity tests performed in this study showed that the post-culture liquids of *B. adusta* CCBAS 930 are not toxic to the tested *E. coli* and *S. aureus* bacteria. In the case of a phytotoxicity assay, discolored post-culture liquids were characterized by a similar degree of root length inhibition as the initial solutions of the dyes. In contrast, the germination index of *L. sativum* L. seeds was much higher for decolorized post-culture liquids (88.20% - 95.21%) than the initial solutions of anthraquinone dyes (33.87% - 72.34%). Therefore, it can be concluded that no toxic products of their biotransformation were formed during the removal of anthraquinone dyes in immobilized *B. adusta* CCBAS 930 cultures.

5. Conclusion

The results showed that the use of immobilised mycelium of *B. adusta* CCBAS 930 significantly reduces the time of decolorization of anthraquinone dyes. After 7 days decolorization degree of CA, AB129, ABBB and RBBR was 75.50%, 65.08%, 56.57% and 25,64%, respectively. Activity of HRP-like and VP peroxidase systematically increased during immobilized cultures with the maximum between 4 and 7 days, 35.64 - 80.32 and 50.20 - 106.43 U mg⁻¹ protein, respectively. In case of phenolic compounds systematic decrease in content was observed (50-60%). Decolorized post-culture fluids were characterized by lower bio- and phytotoxicity.

6. Acknowledgements

Publication is funded by the Polish National Agency for Academic Exchange under the International Academic Partnerships Programme from the project „Organization of the 9th International Scientific and Technical Conference entitled Environmental Engineering, Photogrammetry, Geoinformatics – Modern Technologies and Development Perspectives”.

7. References

- 1 Q. Huang, R.A. Pinto, K. Griebenow, R., Schweitzer-Stenner, W.J. Weber, J. Am. Chem. Soc. **127** (5), 1431–1437 (2005)
- 2 Y. Fu, T. Viraraghavan, Bioresour Technol **79**, 251–262 (2001)
- 3 R.C. Kuhad, N. Sood, K.K. Tripathi, A. Singh O.P. Adv. Appl. Microbiol. **56**, 185–213 (2004)
- 4 T. Kornilowicz-Kowalska, K. Rybczyńska, K. Cent. Eur. J. Biol. **7**, 948-956 (2012)
- 5 T. Kornilowicz-Kowalska, K. Rybczyńska World J. Microbiol. Biotechnol. **30**, 1725-1736 (2014)
- 6 T. Kornilowicz-Kowalska, H. Iglík, Pol. J. Environ. Stu. **16**, 4720-482 (2007)
- 7 Z. Bouabidi, M. El-Naas, Z. Zhang, Environ. Chem. Lett. **17**, 241-257 (2018)
- 8 M.B. Kurade, T.R. Waghmode, J.Q. Xiong, S.P. Govindwar, B.H. Jeon, 2019. J. Clean. Prod. **213**, 884-891 (2019)
- 9 T. Kornilowicz-Kowalska, M. Wrzosek, G. Ginalska, H. Iglík, R. Bancercz, Enzyme Microb. Technol. **38**, 583-590 (2006)
- 10 K. Rybczyńska-Tkaczyk, A. Święciło, K.A. Szychowki, T. Kornilowicz-Kowalska. Ecotox. Environ. Safte. **147**, 776-787 (2018)
- 11 A. Święciło, K. Rybczyńska-Tkaczyk, A. Najda, A., Krzepińko, R. Prażak, G. Zawiślak, G., LWT Food Sci. Technol. **94**, 96-102 (2018)
- 12 M.M. Bradford, Anal. Biochem. **7**, 248– 254 (1976)
- 13 K. Rybczyńska-Tkaczyk, T. Kornilowicz-Kowalska, Int. Biodeterior. Biodegrad. **117**,131-140 (2017)
- 14 I Osaka, S. Hefty, Antimicrob. Agents Chemother. **57**, 2838 –2840 (2013)
- 15 A. Dominguez, S.R. Couto, M.A. Sanromán, World J. Microbiol. Biotechnol. **21**, 405-409 (2005)
- 16 B.B. Hameed, Z.Z. Ismail, Biochem. Eng. J. **137**, 71-77 (2018)
- 17 E. Forgacs, T. Cserha'ti, G. Oros G Removal of synthetic dyes from wastewaters: a review. Environ Int **30**:953–971

- 18 M.J. Lopez, G. Guisado, M.C. Vargas-Garcia, F. Suarez-Estrella, J. Moreno, *Enzyme Microb. Technol.* **40**, 42–45 (2006)
- 19 T. Kornilłowicz-Kowalska, K. Rybczyńska, *Int. J. Environ. Sci. Technol.* **12**, 2673–2686 (2015)
- 20 R. Tinoco, J. Verdin, R. Vazquez-Duhalt, J. *Mol. Cat. B: Enzym.* **46**, 1-7 (2007)
- 21 P.R. Moreira, F. Bouillenne, E. Almeida-Vara, F. Xavier Malcata, J.M. Frére, J.C. Duarte, *Enzyme Microb. Technol.* **38**, 28-3 (2006)