

CONTINUED FRACTIONS OF HIGHER ORDER POLYGONAL NUMBERS WITH RESPECT TO ORDER AND RANK

B.Anitha¹, **P.Balamurugan**²

¹*PG and Research Department of Mathematics, National College (Autonomous), (Affiliated to Bharathidasan University), Trichy-620001.*

²*Department of Mathematics, M.Kumarasamy College of Engineering (Autonomous), Karur – 639113.*

Abstract: Developing number sequence based on polygonal numbers is an enthusiastic field in number theory. As tetrahedral numbers are similar to pyramids, one of the Seven wonders of the World, yields a unique copiousness in its suitability. In number theory study of pyramidal numbers vary in richness and variety. Also the study of continued fractions is a fastly developing field.

Keywords: *continued fraction, pyramidal number, order, rank.*

Notations

1. $\langle p_0, p_1, p_2, p_3, \dots, p_n \rangle$ – continued fraction expansion
2. $PN(n, r)$ – pyramidal number of order 'n' and rank 'r'
3. $PN(3, r)$ – triangular pyramidal number
4. $PN(4, r)$ – square pyramidal number

1. Introduction

Representation of rational numbers as continued fractions is continuously studied by various mathematicians. The convergence of continued fractions is specially studied and applied in finding the best rational gear ratio approximations. Higher dimensional polygonal numbers namely pyramidal numbers combined with continued fractions is really mind boggling. Finding solution of cubic equations with figurate numbers as coefficients in terms of continued fractions is found in. Centered polygonal numbers taken in various forms as continued fraction as given in [3,9] is really a motivating factor for this paper.

1Correspondingauthor: anithamaths2010@gmail.com

2. Pyramidal number

The polygonal number of higher order namely, pyramidal number of order n and rank r is represented by the formula

$$PN(n, r) = \frac{n(n+1)(n(r-2) - (r-5))}{6}$$

2.1 Continued Fraction

An expression of the form

$$\frac{a}{b} = p_0 + \frac{q_0}{p_1 + \frac{q_1}{p_2 + \frac{q_2}{p_3 + \dots}}}$$

where p_i, q_i are real or complex numbers is called a continued fraction.

2.1 Theorem

If ' n ' and ' r ' are order and rank of the corresponding pyramidal numbers then the continued fraction of $\frac{PN(n,r)}{PN(n+1,r)}$ is given by

$$\frac{PN(n,r)}{PN(n+1,r)} = \left\{ \begin{array}{l} \langle 0, 1, 2k-1, \frac{(3k-1)r - 3(2k-1)}{k} \rangle, \text{ if } n = 6k-2 \\ \langle 0, 1, 2k-1, 1, \frac{(6k-1)r - (6k-5)}{2r(6k-1) - (4(6k-1) + 3(2k-1))} \rangle, \text{ if } n = 6k-1 \\ \langle 0, 1, 2k-1, 1, \frac{2kr - (3k-1)}{k(r-3)} \rangle, \text{ if } n = 6k \\ \langle 0, 1, 2k, \frac{(6k+1)r - 12k}{2k+1} \rangle, \text{ if } n = 6k+1 \\ \langle 0, 1, 2k, 1, \frac{(3k+1)r - (3k-1)}{(6k+2)r - (15k+4)} \rangle, \text{ if } n = 6k+2 \\ \langle 0, 1, 2k, 1, \frac{(4k+2)r - (6k+1)}{(2k+1)(r-3)} \rangle, \text{ if } n = 6k+3, k = 1,2,3,4.. \end{array} \right.$$

Proof:

$$\begin{aligned} \frac{PN(n,r)}{PN(n+1,r)} &= \frac{n(n+1)(n(r-2) - (r-5))/6}{(n+1)(n+2)((n+1)(r-2) - (r-5))/6} \\ &= \frac{n(nr - 2n - r + 5)}{(n+2)(nr + r - 2n - 2 - r + 5)} \\ &= \frac{n(nr - 2n - r + 5)}{(n+2)(nr - 2n + 3)} \end{aligned}$$

$$= \frac{n^2r - 2n^2 - nr + 5n}{n^2r - 2n^2 + 3n + 2nr - 4n + 6}$$

$$\frac{PN(n,r)}{PN(n+1,r)} = \frac{n^2r - 2n^2 - nr + 5n}{n^2r - 2n^2 + 2nr - n + 6}$$

Case (i) Take $n = 6k - 2$

$$\frac{PN(6k-2,r)}{PN(6k-1,r)} = \frac{(6k-2)^2r - 2(6k-2)^2 - (6k-2)r + 5(6k-2)}{(6k-2)^2r - 2(6k-2)^2 + 2r(6k-2) - (6k-2) + 6}$$

$$= \frac{36k^2r + 4r - 24kr - 72k^2 - 8 + 48k - 6kr + 2r + 30k - 10}{36k^2r + 4r - 24kr - 72k^2 - 8 + 48k + 12kr - 4r - 6k + 8}$$

$$= \frac{36k^2r - 72k^2 - 30kr + 6r + 78k - 18}{36k^2r - 72k^2 - 12kr + 42k}$$

$$= 0 + \frac{1}{\frac{36k^2r - 72k^2 - 12kr + 42k}{36k^2r - 72k^2 - 30kr + 78k + 6r - 18}}$$

$$= 0 + \frac{1}{1 + \frac{1}{\frac{36k^2r - 72k^2 - 30kr + 78k + 6r - 18}{18kr - 36k - 6r + 18}}}$$

$$= 0 + \frac{1}{1 + \frac{1}{(2k-1) + \frac{6k}{18kr - 36k - 6r + 18}}}$$

$$= 0 + \frac{1}{1 + \frac{1}{(2k-1) + \frac{1}{\frac{1}{(3k-1)r - 3(2k-1)}}}}$$

$$\frac{PN(6k-2,r)}{PN(6k-1,r)} = \langle 0, 1, 2k - 1, \frac{(3k-1)r - 3(2k-1)}{k} \rangle$$

2.2 Illustrations

Value	Ratios	Continued fraction
$k = 1$	$\frac{PN(4,r)}{PN(5,r)}$	$\langle 0, 1, 1, \frac{2r-3}{1} \rangle$
$k = 2$	$\frac{PN(10,r)}{PN(11,r)}$	$\langle 0, 1, 3, \frac{5r-9}{2} \rangle$

Case (ii) Take $n = 6k - 1$

$$\frac{PN(6k-1,r)}{PN(6k,r)} = \frac{(6k-1)^2r - 2(6k-1)^2 - (6k-1)r + 5(6k-1)}{(6k-1)^2r - 2(6k-1)^2 + 2r(6k-1) - (6k-1) + 6}$$

$$\begin{aligned}
 &= \frac{(36k^2 + 1 - 12k)r - 2(36k^2 + 1 - 12k) - 6kr + r + 30k - 5}{(36k^2 + 1 - 12k)r - 2(36k^2 + 1 - 12k) + 12kr - 2r - 6k + 1 + 6} \\
 &= \frac{36k^2r + r - 12kr - 72k^2 - 2 + 24k - 6kr + r + 30k - 5}{36k^2r + r - 12kr - 72k^2 - 2 + 24k + 12kr - 2r - 6k + 7} \\
 &= \frac{36k^2r - 72k^2 - 18kr + 2r + 54k - 7}{36k^2r - 72k^2 - r + 18k + 5} \\
 &= 0 + \frac{1}{\frac{36k^2r - 72k^2 - r + 18k + 5}{36k^2r - 72k^2 - 18kr + 2r + 54k - 7}} \\
 &= 0 + \frac{1}{1 + \frac{18kr - 3r - 36k + 12}{36k^2r - 72k^2 - 18kr + 2r + 54k - 7}} \\
 &= 0 + \frac{1}{1 + \frac{1}{\frac{36k^2r - 72k^2 - 18kr + 2r + 54k - 7}{18kr - 3r - 36k + 12}}} \\
 &= 0 + \frac{1}{1 + \frac{1}{(2k-1) + \frac{6kr - r - 6k + 5}{18kr - 3r - 36k + 12}}} \\
 &= 0 + \frac{1}{1 + \frac{1}{(2k-1) + \frac{1}{\frac{18kr - 3r - 36k + 12}{6kr - r - 6k + 5}}}} \\
 &= 0 + \frac{1}{1 + \frac{1}{(2k-1) + \frac{1}{1 + \frac{6kr - r - 6k + 5}{12kr - 2r - 30k + 7}}}}}
 \end{aligned}$$

$$\frac{PN(6k - 1, r)}{PN(6k, r)} = \langle 0, 1, 2k - 1, 1, \frac{(6k - 1)r - (6k - 5)}{2r(6k - 1) - (4(6k - 1) + 3(2k - 1))} \rangle$$

2.3 Illustrations:

Value	Ratios	Continued fraction
$k = 1$	$\frac{PN(5, r)}{PN(6, r)}$	$\langle 0, 1, 1, 1, \frac{5r - 1}{10r - 23} \rangle$
$k = 2$	$\frac{PN(11, r)}{PN(12, r)}$	$\langle 0, 1, 3, 1, \frac{11r - 7}{22r - 53} \rangle$

Case (iii) Take $n = 6k$

$$\begin{aligned}
 \frac{PN(6k, r)}{PN(6k + 1, r)} &= \frac{(6k)^2r - 2(6k)^2 - 6kr + 5(6k)}{(6k)^2r - 2(6k)^2 + 2r(6k) - 6k + 6} \\
 &= \frac{36k^2r - 72k^2 - 6kr + 30k}{36k^2 - 72k^2 + 12kr - 6k + 6}
 \end{aligned}$$

$$\begin{aligned}
 &= 0 + \frac{1}{\frac{36k^2 - 72k^2 + 12kr - 6k + 6}{36k^2r - 72k^2 - 6kr + 30k}} \\
 &= 0 + \frac{1}{1 + \frac{18kr - 36k + 6}{36k^2r - 72k^2 - 6kr + 30k}} \\
 &= 0 + \frac{1}{1 + \frac{1}{\frac{36k^2r - 72k^2 - 6kr + 30k}{18kr - 36k + 6}}} \\
 &= 0 + \frac{1}{1 + \frac{1}{(2k-1) + \frac{12kr - 18k + 6}{18kr - 36k + 6}}} \\
 &= 0 + \frac{1}{1 + \frac{1}{(2k-1) + \frac{1}{\frac{3kr - 6k + 1}{2kr - 3k + 1}}}} \\
 &= 0 + \frac{1}{1 + \frac{1}{(2k-1) + \frac{1}{1 + \frac{kr - 3k}{2kr - 3k + 1}}}} \\
 &= 0 + \frac{1}{1 + \frac{1}{(2k-1) + \frac{1}{1 + \frac{1}{\frac{k(r-3)}{k(r-3)}}}}}
 \end{aligned}$$

$$\frac{PN(6k, r)}{PN(6k + 1, r)} = \langle 0, 1, 2k - 1, 1, \frac{2kr - (3k - 1)}{k(r - 3)} \rangle$$

2.4 Illustrations

Value	Ratios	Continued fraction
$k = 1$	$\frac{PN(6, r)}{PN(7, r)}$	$\langle 0, 1, 1, 1, \frac{2r - 2}{1(r - 3)} \rangle$
$k = 2$	$\frac{PN(12, r)}{PN(13, r)}$	$\langle 0, 1, 3, 1, \frac{4r - 5}{2(r - 3)} \rangle$

Case (iv) Take $n = 6k + 1$

$$\begin{aligned}
 \frac{PN(6k + 1, r)}{PN(6k + 2, r)} &= \frac{(6k + 1)^2r - 2(6k + 1)^2 - (6k + 1)r + 5(6k + 1)}{(6k + 1)^2r - 2(6k + 1)^2 + 2r(6k + 1) - (6k + 1) + 6} \\
 &= \frac{36k^2r + r + 12kr - 72k^2 - 2 - 24k - 6kr - r + 30k + 5}{36k^2r + r + 12kr - 72k^2 - 2 - 24k + 12kr + 2r - 6k + 5} \\
 &= \frac{36k^2r - 72k^2 + 6kr + 6k + 3}{36k^2r - 72k^2 + 24kr + 3r - 30k + 3}
 \end{aligned}$$

$$\begin{aligned}
 &= 0 + \frac{1}{1 + \frac{1}{\frac{36k^2r - 72k^2 + 6kr + 6k + 3}{3r + 18kr - 36k}}} \\
 &= 0 + \frac{1}{1 + \frac{1}{2k + \frac{18kr - 36k + 3r}{6k + 3}}} \\
 &= 0 + \frac{1}{1 + \frac{1}{2k + \frac{(6k + 1)r - 12k}{2k + 1}}} \\
 \frac{PN(6k + 1, r)}{PN(6k + 2, r)} &= \langle 0, 1, 2k, \frac{(6k + 1)r - 12k}{2k + 1} \rangle
 \end{aligned}$$

2.5 Illustrations :

Value	Ratios	Continued fraction
$k = 1$	$\frac{PN(7, r)}{PN(8, r)}$	$\langle 0, 1, 2, \frac{7r - 12}{3} \rangle$
$k = 2$	$\frac{PN(13, r)}{PN(14, r)}$	$\langle 0, 1, 4, \frac{13r - 24}{5} \rangle$

Case (v) Take $n = 6k + 2$

$$\begin{aligned}
 \frac{PN(6k + 2, r)}{PN(6k + 3, r)} &= \frac{(6k + 2)^2r - 2(6k + 2)^2 - (6k + 2)r + 5(6k + 2)}{(6k + 2)^2r - 2(6k + 2)^2 + 2r(6k + 2) - (6k + 2) + 6} \\
 &= \frac{(36k^2 + 4 + 24k)r - 2(36k^2 + 4 + 24k) - 6kr - 2r + 30k + 10}{(36k^2 + 4 + 24k)r - 2(36k^2 + 4 + 24k) + 12kr + 4r - 6k - 2 + 6} \\
 &= 0 + \frac{1}{\frac{36k^2r - 72k^2 + 36kr + 8r - 54k - 4}{36k^2r - 72k^2 + 18kr + 2r - 18k + 2}} \\
 &= 0 + \frac{1}{1 + \frac{18kr + 6r - 36k - 6}{36k^2r - 72k^2 + 18kr + 2r - 18k + 2}} \\
 &= 0 + \frac{1}{1 + \frac{1}{\frac{36k^2r - 72k^2 + 18kr + 2r - 18k + 2}{18kr + 6r - 36k - 6}}} \\
 &= 0 + \frac{1}{1 + \frac{1}{2k + \frac{6kr + 2r - 6k + 2}{18kr + 6r - 36k - 6}}} \\
 &= 0 + \frac{1}{1 + \frac{1}{2k + \frac{1}{1 + \frac{6kr + 2r - 6k + 2}{12kr + 4r - 30k - 8}}}}
 \end{aligned}$$

$$= 0 + \frac{1}{1 + \frac{1}{2k + \frac{1}{1 + \frac{1}{1 + \frac{3kr+r-3k+1}{6kr+2r-15k-4}}}}}$$

$$\frac{PN(6k + 2, r)}{PN(6k + 3, r)} = \langle 0, 1, 2k, 1, \frac{(3k + 1)r - (3k - 1)}{(6k + 2)r - (15k + 4)} \rangle$$

2.6 Illustrations :

Value	Ratios	Continued fraction
$k = 1$	$\frac{PN(8, r)}{PN(9, r)}$	$\langle 0, 1, 2, 1, \frac{4r - 2}{8r - 19} \rangle$
$k = 2$	$\frac{PN(14, r)}{PN(15, r)}$	$\langle 0, 1, 4, 1, \frac{7r - 5}{14r - 34} \rangle$

(vi) Take $n = 6k + 3$

$$\begin{aligned} \frac{PN(6k + 3, r)}{PN(6k + 4, r)} &= \frac{(6k + 3)^2r - 2(6k + 3)^2 - (6k + 3)r + 5(6k + 3)}{(6k + 3)^2r - 2(6k + 3)^2 + 2r(6k + 3) - (6k + 3) + 6} \\ &= \frac{36k^2r - 72k^2 + 30kr + 6r - 42k - 3}{36k^2r - 72k^2 + 48kr + 15r - 78k - 15} \\ &= 0 + \frac{1}{\frac{36k^2r - 72k^2 + 48kr + 15r - 78k - 15}{36k^2r - 72k^2 + 30kr + 6r - 42k - 3}} \\ &= 0 + \frac{1}{1 + \frac{18kr + 9r - 36k - 12}{36k^2r - 72k^2 + 30kr + 6r - 42k - 3}} \\ &= 0 + \frac{1}{1 + \frac{1}{\frac{36k^2r - 72k^2 + 30kr + 6r - 42k - 3}{18kr + 9r - 36k - 12}}} \\ &= 0 + \frac{1}{1 + \frac{1}{2k + \frac{18kr + 9r - 36k - 12}{12kr + 6r - 18k - 3}}} \\ &= 0 + \frac{1}{1 + \frac{1}{2k + \frac{1}{1 + \frac{12kr + 6r - 18k - 3}{6kr + 3r - 18k - 9}}}} \\ &= 0 + \frac{1}{1 + \frac{1}{2k + \frac{1}{1 + \frac{4kr + 2r - 6k - 1}{2kr + r - 6k - 3}}}} \end{aligned}$$

$$\frac{PN(6k + 3, r)}{PN(6k + 4, r)} = \langle 0, 1, 2k, 1, \frac{(4k + 2)r - (6k + 1)}{(2k + 1)(r - 3)} \rangle$$

2.7 Illustrations :

Value	Ratios	Continued fraction
$k = 1$	$\frac{PN(9, r)}{PN(10, r)}$	$\langle 0, 1, 2, 1, \frac{6r - 7}{3(r - 3)} \rangle$
$k = 2$	$\frac{PN(15, r)}{PN(16, r)}$	$\langle 0, 1, 4, 1, \frac{10r - 13}{5(r - 3)} \rangle$

2.8 Theorem :

If 'n' and 'r' are order and rank of the corresponding pyramidal numbers then the continued function of

$$\frac{PN(n, r)}{PN(n, r + 1)} = \langle 0, 1, (r - 2), \frac{n - 1}{3} \rangle$$

Proof:

$$\begin{aligned} \frac{PN(n, r)}{PN(n, r + 1)} &= \frac{(n(n + 1)(n(r - 2) - (r - 5))) / 6}{(n(n + 1)(n(r + 1 - 2) - (r + 1 - 5))) / 6} \\ &= 0 + \frac{1}{\frac{nr - n - r + 4}{nr - 2n - r + 5}} \\ &= 0 + \frac{1}{1 + \frac{1}{(r - 2) + \frac{1}{\frac{n - 1}{3}}}} \end{aligned}$$

$$\frac{PN(n, r)}{PN(n, r + 1)} = \langle 0, 1, (r - 2), \frac{n - 1}{3} \rangle$$

2.9 Illustrations

2.10

Ratios	Continued fraction
$\frac{PN(n, 3)}{PN(n, 4)}$	$\langle 0, 1, 1, \frac{n - 1}{3} \rangle$
$\frac{PN(n, 5)}{PN(n, 6)}$	$\langle 0, 1, 3, \frac{n - 1}{3} \rangle$

3. Conclusion

Continued fractions of pyramidal numbers with higher orders in the numerator are studied here. When the ratios of continued fractions with consecutive ranks are taken six cases arise whenever for consecutive ordered fractions only a single case arise. Consecutive ordered pyramidal numbers have been taken up for study.

References

- [1] A.Gnanam, S.Krithika,“ Ratios of Polygonal Numbers as Continued Fractions ”, International journal of Engineering Science, Advanced Computing and Bio Technology, Vol.8,No.3, pages 143-155, July-September 2017.
- [2] P.Balamurugan, A.Gnanam,“Pattern classification of continued fractions with triangular number as base”, International journal of pure and applied mathematics, Vol 119,No 15, Pages 3413-3418, 2018.
- [3] P.Balamurugan, A.Gnanam,“Pattern classification of continued fractions with centered polygonal number as base”, International journal of Management. , IT and Engineering, Vol 9, Issue 1,Pages 156-162 (Special issue HICAAMMC) -January-2019.
- [4] P.Balamurugan, A.Gnanam, B.Anitha,“ Expression of ratios of polygonal numbers as continued fractions”, Advances and Applications in Mathematical Sciences, Vol 18, Issue 10, Pages 997-1006, August 2019.
- [5] B.Anitha,“ Expression of ratios of pentatope numbers as continued fractions”, Advances and Applications in Mathematical Sciences, Vol 21, Issue 7,Pages 3873-3883,May 2022.
- [6] B.Anitha,“ Some remarkable observations of farey sequence”, Advances and Applications in Mathematical Sciences,Vol 21, Issue 7,Pages 3885-3893,May 2022.
- [7] A.Venkatachalam, P.Balamurugan,“Pattern classification of continued fractions with square number as base”,Journal of Physics: Conference Series, Vol 1362, Issue 1, pp 0120843, 2019.
- [8] A.Gnanam, B.Anitha,, “ Sums of squares of Fibonacci numbers with prime indices”, Journal of Applied Mathematics and Physics”, Vol 3, Issue 12 ,Pages 1619-1623,2015.
- [9] P.Balamurugan, A.Gnanam, R.Senthilkumar, Sivaranjani M,“Continued fractions of ratios of polygonal numbers with consecutive orders and ranks”, International Journal of Advanced Science and Technology,.Vol 29,issue 7s,Pages 1550 - 1556.
- [10] S.Krithika, A.Gnanam,“Solution of a cubic Equation with triangular number as coefficients ”, International journal of Recent Technology and Engineering, Vol.8, Issue 3, pages 8867-8870, 2019.
- [11] A.Gnanam, B.Anitha,, “ Sums of squares of polygonal numbers”,Advances in Pure Mathematics”, Vol 6, Issue 4 ,Pages 297-301, August 2016.